

# Triple Antibiotic Paste Instructions



## Your Order Contains 3 Kits

**NOTE:** While the triple antibiotic paste can be used for up to 5 days after mixing, it is recommended that it be mixed immediately before use.

## What is supplied in each kit:

- (1) 6 cc Luer lock syringe with triple antibiotic powder and syringe
- (1) 6 cc Luer lock syringe with PEG-PG liquid and syringe cap
- (1) Female-to-female Luer-lock connector
- (1) 1.2cc Luer lock syringe with cap
- (2) Pre-bent disposable flow tips



6 cc Luer Lock Syringe with Triple Antibiotic Powder (1g)



6 cc Luer lock syringe with PEG-PG liquid (1mL)



Female-to-Female Luer Lock Connector



1.2cc Luer Lock Syringe with Cap



Pre-Bent Disposable Flow Tips

## Component Assembly and Mixing Instructions

**Important:** Wash hands thoroughly with soap and water, dry hands, and wear gloves before mixing medication.


- 1. Syringe with triple antibiotic powder:** Hold the syringe upright (the syringe cap should be facing up). Tap the syringe a few times on a flat surface to dislodge any powder that may have attached to the syringe cap.
- 2.** Remove the cap from the syringe by twisting it off counter-clockwise.
- 3.** Attach the female-to-female Luer lock connector to the syringe by twisting it clockwise.
- 4. Syringe with PEG-PG liquid:** Hold the syringe upright (the syringe cap should be facing up).
- 5.** Remove the cap from the syringe by twisting it off counter-clockwise.
- 6.** Attach syringe with PEG-PG liquid to the Luer lock connector that is attached to the syringe with triple antibiotic powder by twisting it on clockwise.

# Triple Antibiotic Paste Instructions



**7. Mixing the syringes:** Mix the powder and liquid by alternately depressing the plungers, first on the liquid syringe and then on the powder syringe for ten mixing strokes. Each mixing stroke is one complete compression of the gel syringe plunger followed by one complete compression of the powder syringe plunger.

**Note:** if you find that depressing the plungers is difficult, you may use a tabletop to assist with mixing. While holding one syringe, place the plunger against the tabletop and push down the syringe body. This will push the contents into the other syringe. Now repeat with the other syringe until mixed.



## Storage

- Store unprepared syringes at room temperature.
- Store activated medication in refrigerator.

## Expiration Date

- Unprepared medication has an expiration date of 6 months from the date compounded. Do not use beyond expiration date.
- Once medication is activated (once the syringes are mixed) the **expiration date is 5 days**. Do not use beyond expiration date.